

UGC POINT ACADEMY


LEADING INSTITUTE FOR CSIR-NET/JRF, GATE & JAM

TATA INSTITUTE OF FUNDAMENTAL RESEARCH PHYSICS DECEMBER-2012

1. The different 2×2 matrices A and B are found to have the same eigen-values. It is then correct to state that $A = SBS^{-1}$ where S can be a
- (a) traceless 2×2 matrix
(b) Hermitian 2×2 matrix
(c) unitary 2×2 matrix
(d) arbitrary 2×2 matrix

2. The function $f(x)$ represents the nearest integer less than x , e.g.
 $f(3.14) = 3$
The derivative of this function (for arbitrary x) will be given in terms of the integers n as $f'(x) =$
- (a) 0
(b) $\sum_n \delta(x-n)$
(c) $\sum_n |x-n|$
(d) $\sum_n f(x-n)$

3. Two masses M_1 and M_2 ($M_1 < M_2$) are suspended from a perfectly rigid horizontal support by a system of three taut mass-less wires W_1 , W_2 and W_3 , as shown in the figure. All the three wires have identical cross-section and elastic properties and are known to be very strong.



If the mass M_2 is increased gradually, but without limit, we should expect the wires to break in the following order:

- (a) first W_2 , then W_1
(b) first W_1 , then W_2
(c) first W_2 , then W_3
(d) first W_3
4. A high-velocity missile, travelling in a horizontal line with a kinetic energy of 3.0 Giga-Joules (GJ), explodes in flight and breaks into two pieces A and B of equal mass. One of these pieces (A) flies off in a straight line perpendicular to the original direction in which the missile was moving and its kinetic energy is found to be 2.0 GJ. If gravity can be neglected for such high-velocity projectiles, it follows that the other piece (B) flew off in a direction at an angle with the original direction of
- (a) 30°
(b) $33^\circ 24'$
(c) 45°
(d) 60°
5. Consider a spherical planet, rotating about an axis passing through its centre. The velocity of a point on its equator is v_{eq} . If the acceleration due to gravity g measured at the equator is half of the value of g measured at one of the poles, then the escape velocity for a particle shot upwards from the pole will be
- (a) $\frac{v_{eq}}{2}$
(b) $\frac{v_{eq}}{\sqrt{2}}$
(c) $\sqrt{2}v_{eq}$
(d) $2v_{eq}$

27-G, 2nd Floor, Jia Sarai, Near IIT, Hauz Khas New Delhi-110016

Tel: 011-26521410, 26855515 Mob: 09654680505, 07503646974

E-Mail: info@ugcpoint.in Website: www.ugcpoint.in


6. A dynamical system with two degrees of freedom, has generalized coordinates q_1 and q_2 , and kinetic energy

$$T = \lambda \dot{q}_1 \dot{q}_2$$

If the potential energy is $V(q_1, q_2) = 0$, the correct form of the Hamiltonian for this system is


- (a) $\frac{P_1 P_2}{\lambda}$ (b) $\lambda \dot{q}_1 \dot{q}_2$ (c) $\frac{(P_1 \dot{q}_1 + P_2 \dot{q}_2)}{2}$ (d) $\frac{(P_1 q_2 + P_2 q_1)}{2}$

7. An ideal liquid of density 1 gm/cc is flowing at a rate of 10 gm/s through a tube with varying cross-section, as shown in the figure



Two pressure gauges attached at the points A and B (see figure) show readings of P_A and P_B respectively. If the radius of the tube at the points A and B is 0.2 m and 1.0 cm respectively, then the difference in pressure ($P_B - P_A$), in units of dyne cm^{-2} is closet to

- (a) 100 (b) 120 (c) 140 (d) 160
8. Unpolarised light of intensity I_0 passes successively through two identical linear polarisers A and B, placed such that their polarization axes are at an angle of 45° (see figure) with respect to one another.



Assuming A and B to be perfect polarisers (i.e. no absorption losses), the intensity of the transmitted light will be $I_T =$

- (a) $\frac{I_0}{4}$ (b) $\frac{I_0}{2\sqrt{2}}$ (c) $\frac{I_0}{2}$ (d) $\frac{I_0}{\sqrt{2}}$


9. Three equal charges Q are successively brought from infinity and each is placed at one of the three vertices of an equilateral triangle. Assuming the rest of the universe as a whole to be neutral, the energy E_0 of the electrostatic field will increase, successively to

$$E_0 + \Delta_1, \quad E_0 + \Delta_1 + \Delta_2, \quad E_0 + \Delta_1 + \Delta_2 + \Delta_3$$

Where $\Delta_1 : \Delta_2 : \Delta_3 =$


- (a) 1:2:3 (b) 1:1:1 (c) 0:1:1 (d) 0:1:2

10. Five sides of a hollow metallic cube are grounded and the sixth side is insulated from the rest and is held at a potential Φ (see figure)



The potential at the centre O of the cube is

- (a) 0 (b) $\frac{\Phi}{6}$ (c) $\frac{\Phi}{5}$ (d) $\frac{2\Phi}{3}$
11. Consider a sealed but thermally container of total volume V, which is in equilibrium with a thermal bath at temperature T. The container is divided into two equal chambers by a thin but impermeable partition. One of these chambers contains an ideal gas, while the other half is a vacuum (see figure)



If the partition is removed and the ideal gas is allowed to expand and fill the entire container, then the entropy per molecule of the system will increase by an amount


- (a) $2k_B$ (b) $k_B \ln(1/2)$ (c) $k_B \ln 2$ (d) $\frac{(k_B \ln 2)}{2}$
12. When a gas is enclosed in an impermeable box and heated to a high temperature T, some of the neutral atoms lose an electron and become ions. If the number density of neutral atoms, ions and electrons is N_a, N_i and N_e , respectively, these can be related to the average volume V_a occupied by an atom / ion and the ionization energy E by the relation.
- (a) $N_e(N_a + N_i) = (N_a / V_a) \exp(-E / k_B T)$
 (b) $N_a(N_e / N_i) = (N_a / V_a) \exp(-E / k_B T)$
 (c) $N_e N_i = (N_a / V_a) \exp(+E / k_B T)$
 (d) $N_e N_i = (N_a / V_a) \exp(-E / k_B T)$
13. In a scanning tunneling microscope, a fine platinum needle is held close to a metallic surface in vacuum and electrons are allowed to tunnel across the tiny gap δ between the surface and the needle. The tunneling current I is related to the gap δ , through positive constants a and b, as
- (a) $I = a - b\delta$ (b) $I = a + b\delta$ (c) $\log I = a - b\delta$ (d) $\log I = a + b\delta$

14. A particle in a one-dimensional potential has the wave-function

$$\Psi(x) = \frac{1}{\sqrt{a}} \exp\left(-\frac{|x|}{a}\right)$$


Where a is a constant. It follows that for a positive constant V_0 , the potential $V(x) =$

- (a) $V_0 x^2$ (b) $V_0 |x|$ (c) $-V_0 \delta(x)$ (d) $-V_0 / |x|$
15. Consider the high excited states of a hydrogen atom corresponding to large values of the principal quantum number ($n \gg 1$). The wavelength λ of a photon emitted due to an electron undergoing a transition between two such states with consecutive values of n (i.e. $\Psi_{n+1} \rightarrow \Psi_n$) is related to the wavelength λ_α of the K_α line of hydrogen by
- (a) $\lambda = n^3 \lambda_\alpha / 8$ (b) $\lambda = 3n^3 \lambda_\alpha / 8$
(c) $\lambda = n^2 \lambda_\alpha$ (d) $\lambda = 4\lambda_\alpha / n^2$
16. A proton is accelerated to a high E and shot at a nucleus of oxygen ($^{16}_8\text{O}$). In order to penetrate the coulomb barrier and reach the surface of the oxygen nucleus, E must be at least
- (a) 3.6 MeV (b) 1.8 MeV (c) 45 keV (d) 180 eV
17. A monochromatic beam of X-rays with wavelength λ is incident at an angle θ on a crystal with lattice spacing's a and b as sketched in the figure below.



A condition for there to be a maximum in the diffracted X-ray intensity is


- (a) $2\sqrt{a^2 + b^2} \sin \theta = \lambda$ (b) $2b \cos \theta = \lambda$
(c) $2a \cos \theta = \lambda$ (d) $(a + b) \sin \theta = \lambda$
18. Suppose the energy band diagram of a certain pure crystalline solid is as shown in the figure below, where the energy (E) varies with crystal momentum (k) as $E \propto k^2$.



At finite temperature the bottom of the conduction band (CB) is partially filled with electrons (e) and the top of the valence band (VB) is partially filled with holes (h). If an electric field is applied to this solid, both e and h will start moving. If the time between collisions is the same for both e and h , then

- (a) e and h will move with the same speed in opposite directions
- (b) h will on an average achieve higher speed than e
- (c) e will on an average achieve higher speed than h
- (d) e and h will recombine and after a while there will be no flow of charges


19. Consider the circuit shown below




The minimum number of NAND gates required to design this circuit is

- (a) 6
- (b) 5
- (c) 4
- (d) 3





20. Consider the following circuit



If the waveform given below is fed in at V_{in}



Then the waveform at the output V_{out} will be



21. Consider the integral

$$\int_{-p^2}^{+p^2} \frac{dx}{\sqrt{x^2 - p^2}}$$

where p is a constant. The integral has a real nonsingular value if

- (a) $p < -1$ (b) $p > 1$ (c) $p = 1$ (d) $p \rightarrow 0$ (e) $p \rightarrow \infty$

22. If we model the electrons as a uniform sphere of radius r_e , spinning uniformly about an axis passing through its centre with angular momentum $L_e = \frac{\hbar}{2}$, and demand that the velocity of rotation at the equator cannot exceed the velocity c of light in vacuum, then the minimum value of r_e is

- (a) 19.2 fm (b) 0.192 fm (c) 4.8 fm (d) 1960 fm (e) 480 fm


23. The intensity of light coming from a distant star is measured using two identical instruments A and B, where A is placed in a satellite outside the Earth's atmosphere, and B is placed on the Earth's surface. The results are as follows:

| colour | wavelength (nm) | intensity at A (nanoWatts) | intensity at B (nanoWatts) |
|--------|-----------------|----------------------------|----------------------------|
| green | 500 | 100 | 50 |
| red | 700 | 200 | x |

Assuming that there is scattering, but no absorption of light in the Earth's atmosphere at these wavelengths, the value of x can be estimated as

- (a) 137 (b) 147 (c) 157 (d) 167 (e) 177

24. Consider three identical infinite straight wires A, B and C arranged in parallel on a plane as shown in the figure




The wires carry equal currents I with directions as shown in the figure and have mass per unit length m . If the wires A and C are held fixed and the wire B is displaced by a small distance x from its position, then it (B) will execute simple harmonic motion with a time period.

- (a) $2\pi \sqrt{\frac{\pi}{\pi\mu_0}} \left(\frac{d}{I}\right)$ (b) $2\pi \sqrt{\frac{2\pi m}{\mu_0}} \left(\frac{d}{I}\right)$ (c) $2\pi \sqrt{\frac{\pi m}{\mu_0}} \left(\frac{d}{I}\right)$
 (d) $2\pi \sqrt{\frac{m}{2\pi\mu_0}} \left(\frac{d}{I}\right)$ (e) $2\pi \sqrt{\frac{m}{\mu_0}} \left(\frac{d}{I}\right)$

25. The normalized wavefunctions of a Hydrogen atom are denoted by $\Psi_{n,\ell,m}(\vec{x})$, where n, ℓ and m are respectively, the principal, azimuthal and magnetic quantum numbers respectively. Now consider an electron in the mixed state
- $$\Psi(\vec{x}) = \frac{1}{3} \psi_{1,0,0}(\vec{x}) + \frac{2}{3} \psi_{2,1,0}(\vec{x}) + \frac{2}{3} \psi_{3,2,-2}(\vec{x})$$
- The expectation value $\langle E \rangle$ of the energy of this electron, in electron-Volts (eV) will be approximately
- (a) -1.5 (b) -3.7 (c) -13.6 (d) -80.1 (e) +13.6
26. The strongest three lines in the emission spectrum of an interstellar gas cloud are found to have wavelength $\lambda_0, 2\lambda_0$ and $6\lambda_0$ respectively, where λ_0 is a known wavelength. From this we can deduce that the radiating particles in the cloud behave like
- (a) free particles (b) particles in a box (c) harmonic oscillators
(d) rigid rotators (e) hydrogenic atoms
27. When light is emitted from a gas of excited atoms, the lines in the spectrum are Doppler-broadened due to the thermal motion of the emitting atoms. The Doppler width of an emission line of wavelength 500 nanometer (nm) emitted by an excited atom of Argon (${}^{40}_{20}\text{A}$) at room temperature (27°C) can be estimated as
- (a) $5.8 \times 10^{-4} \text{ nm}$ (b) $3.2 \times 10^{-4} \text{ nm}$ (c) $3.2 \times 10^{-3} \text{ nm}$
(d) $2.5 \times 10^{-3} \text{ nm}$ (e) $1.4 \times 10^{-3} \text{ nm}$
28. In a nuclear reactor, Plutonium (${}^{239}_{94}\text{Pu}$) is used as fuel, releasing energy by its fission into isotopes of Barium (${}^{146}_{54}\text{Ba}$) and Strontium (${}^{91}_{38}\text{Sr}$) through the reaction
- $${}^{239}_{94}\text{Pu} + {}^1_0\text{n} \rightarrow {}^{146}_{54}\text{Ba} + {}^{91}_{38}\text{Sr} + 3 \times {}^1_0\text{n}$$
- The binding energy (B.E) per nucleon of each of these nuclides is given in the table below:
- | Nuclide | ${}^{239}_{94}\text{Pu}$ | ${}^{146}_{54}\text{Ba}$ | ${}^{91}_{38}\text{Sr}$ |
|------------------------|--------------------------|--------------------------|-------------------------|
| B.E. per nucleon (MeV) | 7.6 | 8.2 | 8.6 |
- Using this information one can estimate the number of such fission reactions per second in a 100 MW reactor as
- (a) 3.9×10^{18} (b) 7.8×10^{18} (c) 5.2×10^{19}
(d) 5.2×10^{18} (e) 8.9×10^{17}
29. Metallic Copper is known to form cubic crystals and the lattice constant is measured from X-ray diffraction studies to be about 0.36 nm. If the specific gravity of Copper is 8.96 and its atomic weight is 63.5, one can conclude that
- (a) the crystals are of simple cubic type
(b) the crystals are of b.c.c. type
(c) the crystals are a mixture of f.c.c. and b.c.c. types
(d) there is insufficient data to distinguish between the previous options

30. The voltage regulator circuit shown in the figure has been made with a zener diode rated at 15V, 200mW. It is required that the circuit should dissipate 150mW power across the fixed load resistor R_L .



For stable operation of this circuit, the input voltage V_i must have a range

- (a) 17.5V – 20.5V
(b) 15.5V – 20.5V
(c) 15.5V – 22.5V
(d) 17.65V – 22.5V
(e) 15.0V – 22.5V

ugc point